


FACT SHEET



Concussion

Author: Laura O. Morris, PT, NCS

Information about Concussions, Dizziness, and Balance



What is a concussion?

Concussion occurs after sustaining a blow to the head (with or without loss of consciousness). It encompasses many symptoms that impact physical, emotional, and mental well-being. Physical symptoms may include headache, dizziness, imbalance, nausea/vomiting, fatigue, difficulty sleeping, double or blurred vision, and sensitivity to light and sound. Emotional symptoms may include irritability, restlessness, anxiety, depression, mood swings, aggression, or decreased tolerance of stress. Cognitive (thinking) symptoms may include difficulty with short or long term memory, confusion, slowed processing, “fogginess,” and difficulty concentrating. Because of the variety of possible symptoms that can impair your day-to-day function, it is very important to see the appropriate healthcare professionals, including specialized physicians, neuropsychologists, and physical therapists.

A concussion is a problem that often goes undiagnosed because it is not something that can be found with diagnostic imaging, such as an MRI or a CT scan. Post-concussion symptoms occur because of chemical and energy changes that occur in the brain due to trauma. The brain becomes inefficient at carrying out normal physiologic processes which then results in the above symptoms.

What steps should I take if I have had a concussion?

After having a concussion, it is important to limit exertion, both physical and cognitive (thinking) because the brain cannot adequately respond to normal demands. Increased levels of physical exertion too soon, such as return to social activities or sports, or cognitive demand, such as returning to school or work, does not allow the brain to heal, and may cause increased symptoms. Once symptoms have improved, you can slowly begin to resume normal activities as long as there is no significant increase in symptoms.

Physical therapy for those who have had a concussion varies from person to person depending on their symptoms and limitations. The physical therapist will complete a comprehensive evaluation to assess current problems. Physical therapy can play a role in reintroducing physical activity. This is a gradual process because it is important not to overload a system that has been compromised by concussion. You must be relatively symptom free in order to progress activity levels.



1111 North Fairfax Street
Alexandria, VA 22314-1488

Phone: 800-999-2782,


Ext 3237

Fax: 703-706-8578

Email: neuropt@apta.org

www.neuropt.org

Concussion



If a person is dizzy or has difficulty maintaining balance, vestibular rehabilitation may be appropriate. The vestibular system, which includes the inner ear and its connections with the brain, is responsible for sensing head movement and helps to maintain balance. In addition, it is responsible for keeping your eyes focused while the head is moving. The inner ear is a very small and fragile structure located in the skull just behind the ear. Trauma to the head can cause inner ear damage. With specific exercise and management skills, dizziness can be reduced significantly or eliminated, and balance can be improved.

It is important that the recommendations of all medical professionals be followed in order to maximize recovery in the shortest amount of time.



1111 North Fairfax Street
Alexandria, VA 22314-1488

Phone: 800-999-2782,
Ext 3237

Fax: 703-706-8578

Email: neuropt@apta.org

www.neuropt.org